

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 February 2001 (01.02.2001)

PCT

(10) International Publication Number
WO 01/07910 A1


- (51) International Patent Classification⁷: G01N 33/48 (74) Agent: BERLINER, Robert; Fulbright & Jaworski, L.L.P., 29th Floor, 865 S. Figueroa Street, Los Angeles, CA 90017-2576 (US).
- (21) International Application Number: PCT/US00/20009
- (22) International Filing Date: 21 July 2000 (21.07.2000) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/358,504 21 July 1999 (21.07.1999) US
- (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 12th Floor, 1111 Franklin Street, Oakland, CA 94607 (US).
- (72) Inventors: ALLBRITTON, Nancy, L.; 22 Virgil Court, Irvine, CA 92612 (US). SIMS, Christopher, E.; 22 Virgil Court, Irvine, CA 92612 (US). BERNIS, Michael, W.; 31132 Via Colinas, Coto de Caza, CA 92679 (US). MEREDITH, Gavin, D.; 2160 Sea Village Circle, Cardiff-by-the-Sea, CA 92007-1221 (US). KRASIEVA, Tatiana, B.; 22 Newton Court, Irvine, CA 92612 (US). TROMBERG, Bruce, J.; 17 Zola Court, Irvine, CA 92612 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD AND APPARATUS FOR DETECTING ENZYMIC ACTIVITY USING MOLECULES THAT CHANGE ELECTROPHORETIC MOBILITY



Best Available Copy

(57) Abstract: The activity of intracellular chemical reactions of molecules is measured by the use of fluorescently labeled substrate molecules (114) that undergo a change in electrophoretic mobility upon chemical reaction such as that catalyzed by an enzyme (112). Specificity is achieved by using labeled substrate molecules (114) that can be acted upon only by specific enzymes (112). Thus, the activity of a specific enzyme (112) or class of enzymes can be determined. Measurements are made with the intracellular presence of such substrate molecules (114), at some time of interest, typically after exposure of the cell (46) to a stimulus (118) that activates a particular enzymatic pathway. To ensure accuracy, measurements must be made in a timely manner so as to minimize chemical reactions occurring subsequent to the time of interest. Fast controllable laser lysis is used to obtain the contents of a single cell (46) into which reporter substrate molecules (114) have been introduced. The cell contents are then subjected to capillary electrophoresis (22) and enzymatic activity is determined by comparing amounts of substrate molecules (114) to amounts of enzymatically altered substrate molecules (128) which are separated by the electrophoresis and identified by the presence of a fluorescent label.

METHOD AND APPARATUS FOR DETECTING ENZYMATIC ACTIVITY USING MOLECULES THAT CHANGE ELECTROPHORETIC MOBILITY

5 *Related Applications*

The present application is a continuation-in-part application of application Serial No. 09/036,706, filed March 6, 1998, entitled "Fast Controllable Laser Lysis of Cells for Analysis".

Background of Invention

10 1. *Field of the Invention*

The invention relates to the field of micro-analytical chemistry in the areas of cellular biochemical and biomedical analysis, and in particular to a class of chemical compounds and method and apparatus for assaying the internal chemical activity of a single cell or selected cells.

15 2. *Description of the Prior Art*

In biology, the individual cell can be thought of as the fundamental unit of life. Ultimately, the chemical processes that occur within single cells give rise to all of the phenomena that we observe in living organisms. Most intracellular chemical processes are mediated by proteins. Typically these
20 proteins are enzymes that catalytically enhance the rates of specific chemical reactions. Very often, intracellular enzymes participate in cascades of chemical reactions known as signal transduction pathways. Each of these signal transduction pathways is composed of sequentially acting enzymes, frequently of a class known as kinases. These enzymes not only interact with
25 and modify the behavior of other proteins within the same pathway, but also influence the operation of other signal transduction pathways. These interacting cascades of signaling molecules and chemical reaction products form complex networks that ultimately regulate processes of cell growth,

proliferation, quiescence, and programmed cell death.

Inappropriate signaling within a cell can give rise to defects in any of these processes and is implicated as the basis for many forms of cancer. To further complicate matters, individual cells display a high degree of

5 heterogeneity in their internal biochemical signaling. Malignant tumors are typically composed of such a heterogeneous group of cells, and cells within the same tumor often utilize disparate and aberrant growth signaling pathways. This heterogeneity makes it necessary to analyze individual cells to elucidate their unique errors in signaling. Where only a small proportion of

10 cells show aberrant signaling, analysis of an entire population will produce an average signal more reflective of the majority rather than of any individual aberrant cell. Thus, the measurement "averages out" the very errors one seeks to detect.

Often there is a genetic basis, in the form of mutations in genes for

15 signaling proteins, for inappropriate signaling. Mutations in the genes for signaling proteins result in the presence of structurally altered, aberrantly acting, signaling proteins. Currently, there exist methods to analyze the DNA and/or RNA of a single cell to detect the presence of signaling protein mutations and thereby infer the presence of mutant signaling proteins.

20 However, it is not currently possible to predict or understand errors in signaling purely from knowledge of the genes or mutations in the genes that occur within a tumor. For example, the aberrant activity of mutant signaling proteins in one pathway may be counteracted or modified by the activities of other proteins, mutant or normal, involved in other pathways. This is very

25 likely since the interior of a tumor cell is a complex mixture of signaling proteins generally with hundreds to thousands of different kinds present simultaneously and participating in pathways that frequently influence one another. To assess the function of any single kind of a signaling protein, its net activity must be measured from within a single living cell.

30 Further, to completely describe the function of an entire pathway, measurements of the activities of all the signaling proteins involved within a

pathway are needed. Such knowledge would be exceptionally useful in the individualized diagnosis and treatment of diseases that involve faulty intracellular signaling since signaling proteins are often prime targets for therapeutic drug intervention.

5 Further, such a capability would revolutionize the ability to conduct research on basic cellular physiology. As noted, kinases represent an exemplary class of protein enzymes that commonly play a key role in intracellular signaling. In fact, aberrant activation of growth promoting kinases is a general feature of tumor cells. Thus protein kinases are promising targets
10 for cancer therapies and the development of kinase antagonist drugs is an area of intense research. These enzymes catalytically enhance the rate at which a covalent chemical bond is formed between a phosphate group and a molecular substrate molecule, frequently, a different protein. This process is termed "phosphorylation". When the substrate itself is an enzyme, such
15 phosphorylation generally either enhances or suppresses the substrate enzyme's chemical activity.

Kinases generally demonstrate substrate specificity such that the preferred substrates of one kind of kinase are not efficiently phosphorylated by other kinds of kinases. Frequently, the preferred substrates for kinases are
20 different kinases; thus, the inappropriate activity of one kinase can result in changes in the activity of multiple downstream kinases within a signaling pathway.

A general method to measure kinase activity within a cell could, in many instances, yield an abundance of information about an entire signaling
25 pathway. However, in order to make such a measurement strict criteria must be met. To begin with, the number of copies per cell of many enzymatic proteins, such as kinases and kinase substrates, can be as low as 100 to 1,000,000 molecules, or about 150 pM to 1.5 μ M in concentration in a typical mammalian cell with a volume of 1 pl. In terms of moles, this is equivalent to
30 a necessary limit range of approximately 1.7×10^{-18} to 1.7×10^{-22} .

Further, since the concentrations of phosphorylated substrates in cells

change on time scales of the order of seconds, the time resolution of the measurement, from the instant the contents of a cell are obtained to the time that the biochemical reactions are terminated, must be sub-second. Most conventional biochemical assays meet neither the temporal resolution nor the sensitivity limits required for these single cell measurements. The temporal resolution requirement can be met through the use of apparatus described in application serial. no. 09/036,706 filed March 6, 1998, and entitled, "Fast Controllable Laser Lysis of Cells for Analysis", to which this continuation-in-part application is related and which parent application is herein expressly incorporated by reference. The necessary degree of sensitivity can be achieved with traditional capillary electrophoresis (CE) methods. Lacking, until now, has been the molecular means to accurately determine the intracellular activity of one or more kinase species or other enzymes.

Current techniques for kinase measurements can generally be divided into three methodologies. The first method uses the phosphorylation of kinase substrates by cellular extracts to estimate the kinase activity that occurs within intact cells. There are a number of major drawbacks to such a method. Because it is not sensitive, the internal contents of large numbers of cells must be pooled. Since the cells are not synchronous with respect to their activation status, a time averaged level of kinase activity is actually measured. Furthermore, during the time required to generate a cellular extract, a time that may span many seconds to minutes, many chemical reactions continue to proceed. This results in a highly distorted representation of the relative amounts of reactants and products as they occurred within actual cells.

Even beyond these difficulties, it is virtually impossible to reproduce the unique chemical microenvironments that occur within cells; thus, the chemical activity observed in extracts can differ greatly from that which actually occurs within intact cells.

A second methodology relies on labeling the kinase within a cell with

specific antibodies, kinase inhibitors, or fluorescent tags and then observing the cells with a fluorescence microscope. Both inactive and active kinase molecules are labeled by this method, and an attempt is then made to infer the activity level of the kinase from its actual location within a cell. However, it is extremely risky to attempt to determine the state of activation of an enzyme solely from its intracellular location. This is particularly true for a number of kinases that are active both when free in the cell and also when bound to intracellular structures.

A third method, still under development, is the use of a fluorescent indicator to actually measure kinase activity. A similar strategy has worked well for the measurement of various intracellular ion concentrations (i.e. Ca^{2+}), but thus far has not been generally applicable to the measurement of kinase activities. One of the drawbacks of this approach is the need for a supraphysiologic concentration of a reporter fluorophore to produce sufficient signal for ready detection-typically 10 μM to 100 μM of fluorescent indicators are necessary within cells for detection by fluorescence microscopy. This is 10 to 1,000,000 times the concentration of typical physiological kinase substrates. At such high concentrations the fluorescent reporter molecule may behave as an inhibitor or may induce other unintended cellular responses such as activation of aberrant signaling pathways. Another difficulty has been that the viscous, highly concentrated, intracellular environment alters the fluorescent properties of the reporter molecule making the detection unreliable.

Finally, the design of fluorescent, environment-sensitive probes for a specific kinase may not be generally applicable to other kinases. Because a given cell or group of cells may contain a large number of different kinases, this may pose a serious problem. The need for broadly applicable, accurate, and sensitive cellular measurement technologies is great.

What is needed is an alternative, yet complementary, strategy to the above approaches. Further, this new approach should be one that it is not limited to the measurement of kinases, but is broadly adaptable to the

measurement of many if not all other kinds of intracellular enzymatic and chemical activities.

Brief Summary of Invention

5 The present invention is a method for measuring intracellular chemical activity. Typically the molecules of interest are intracellular enzymes, proteins that catalyze biochemical reactions within cells. Catalysis by enzymes results in the alteration of the chemical structure of substrate molecules. The invention comprises the use of substrate molecules that undergo a change in
10 electrophoretic mobility when enzymatically acted upon as intracellular reporters of chemical activity. This strategy provides a general method to detect and quantify enzymatic activity by measuring changes in the structure of substrate molecules. A change in electrophoretic mobility allows for separation of enzymatically altered substrate molecules from unaltered
15 substrate molecules by electrophoresis with highly sensitive detection and quantification methods following separation. Even very minor changes in chemical structure can result in measurable differences in electrophoretic mobility. For example, capillary electrophoresis is routinely used to separate stereoisomeric forms of chemical compounds.

20 Most commonly, substrate molecules are selected to report on the activity of a specific enzyme or class of enzymes. These reporter substrate molecules can occur naturally within a cell, be induced within a cell, such as when under the control of an operon or similar genetic control mechanism, or be introduced into a cell by a variety of established methods including, but not
25 limited to, microinjection, electroporation, optoporation, vesicle fusion, pinocytic loading, or by association with membrane permeant peptides. The reporter substrate molecules may be naturally occurring compounds or synthetically derived. Measurements are made with the intracellular presence of such substrate molecules, at some time of interest, typically after exposure
30 of a cell to a stimulus. At the time of interest, part or all of the contents of a single cell or group of cells are harvested for separation by capillary

electrophoresis. To ensure accuracy of measurement, this process must be achieved in a timely manner so as to minimize chemical reactions occurring subsequent to the time of interest. In practice, this means that if a measurement is to be made to within an accuracy of 1%, then less time
5 should pass, from the time of interest to the time of termination of chemical reactions, than is required for the progress of a reaction to change by 1%. Typically, chemical reactions are terminated by physical disruption as with detergents, turbulent mixing, dilution, or separation of molecules by electrophoresis, or by a combination of these processes. Fast controllable
10 laser lysis is an excellent way to harvest, on a sub-second time scale, the contents of a single cell into which reporter substrate molecules have been introduced.

Reporter molecules specific for particular intracellular reactions can be designed to undergo a predictable change in electrophoretic mobility. These
15 molecules can also be modified to be fluorescent and thus detectable in the minute amounts and low concentrations that occur within the tiny volumes of single cells. In the illustrated embodiments, modified peptides are used as representative reporter molecules. Modified peptides are a particularly useful class of such possible reporter molecules for a number of reasons. Foremost,
20 peptides closely mimic natural, endogenous protein substrates and frequently demonstrate a high degree of specificity in the reactions in which they participate and thus report. This can be understood to be the result, in many cases, of the unique primary sequence structure of these peptides. The particular order of amino acid residues that comprises a given peptide
25 conveys unique chemical properties to the molecule, particularly with regards to being recognized and acted upon as a substrate by a particular enzyme. Typically, enzymes whose natural substrates are proteins will efficiently catalyze only those reactions involving molecules that display a particular order or pattern of amino acid residues.

30 Peptides are extremely versatile compounds when used as reporter substrates. Large, diverse libraries of these compounds can be generated

simply by varying the sequence and number of amino acid residues that comprise a given peptide. Furthermore, modifications of basic peptide structure can be made in a number of ways. For example, D- isomer amino acids can be incorporated into peptides in place of the more commonly found L- isomeric forms. Also, unusual or alternate, non-naturally occurring amino acids can be incorporated into peptides. Examples include homo-arginine and homo-lysine in place of arginine and lysine, respectively. Even the backbone peptide bond may be altered. After modifications such as these have been made, the resultant peptides have proven to serve as substrates for enzymatic reactions.

In the illustrated embodiments, the modified peptides are substrates for enzymes known as kinases that alter the peptides by the addition of a phosphate moiety to a specific amino acid(s) within the peptide. Phosphorylation can be effected *in vitro*, in the absence of a cell, and leads to a demonstrable alteration of the peptides' chemical structure and electrophoretic mobility.

Further, the peptides used in the illustrated embodiments are modified at at least one point, frequently the end by covalent addition of a fluorescein group (or other fluorescent group) to allow detection by laser-induced fluorescence. The fluorescent tags can be added to either the amino or carboxyl termini or to specific amino acid residues by chemical techniques well-known to those of skill in the art. Examples of readily modifiable residues include the sulfhydryl group of cysteine, the amino groups of arginine and lysine, and the carboxyl groups of aspartate and glutamate. Example chemical groups that react with sulfhydryl groups include maleimides, iodoacetamides, alkyl halides, aziridines, and epoxides. Chemical groups that react with amines include succinimydyl esters, sulfonyl halides, isothiocyanates, and aldehydes. Chemical groups that react with carboxyls include carbodiimides, hydrazines, alkyl halides, amines, and diazoalkanes. A wealth of different fluorophores, with reactive groups, are commercially available for covalent linkage to peptides at these groups. Examples of

different fluorophores that are available include fluorescein derivatives such as Oregon Green produced by Molecular Probes (Eugene, OR), rhodamine derivatives such as Texas Red and tetramethylrhodamine, NBD (7-nitrobenz-2-oxa-1,3, diazole) derivatives, coumarins, dansyl derivatives, pyrenes, and the cyanine dyes (Cy-3/Cy-5) produced by Amersham/Pharmacia Biotech (Piscataway, NJ). This diversity allows the preparation of peptides that absorb and emit light at wavelengths ranging from the ultraviolet through the visible and into the infrared parts of the spectrum. This is yet another avenue that can be exploited to produce a large variety of reporter substrate peptides with differing spectral properties.

While illustrative, the invention is not contemplated to be limited to use of modified peptides alone or in its scope to the monitoring of reactions by intracellular kinases alone. It is contemplated that a broader range of intracellular chemical reactions can be monitored by this approach. For example, peptide-based reporter substrates may be designed to report on the activities of phosphatases, glycosylases, acetylases, proteases and caspases, and isomerases. It is also contemplated that molecular reporters may be based on other chemical species commonly found in cells such as nucleic acids, carbohydrates, phospholipids, and entire proteins or may be based on compounds (often polymers) not ordinarily found within cells.

Nucleic-acid based reporter molecules could serve as excellent substrates for nucleases, ligases, polymerases, methylases, demethylases, and nucleotide transferases. Lipid based reporter substrates could serve to monitor lipases. Carbohydrate based reporters could serve to monitor oxidoreductases, glycosylases, hydrolases, lyases, and isomerases. Finally since entire proteins are frequently the natural substrates for intracellular chemical reactions, reporters based on these could prove to yield extremely accurate information about intracellular physiology.

Reporter molecules that comprise entire proteins may be based on the green fluorescent protein (GFP) from the jellyfish *Aequorea victoria*. This protein is intrinsically fluorescent and has been used extensively in basic

biological research. The gene for the protein has been cloned and mutated to produce a diversity of GFP-like fluorescent proteins that have unique absorption and emission spectra. A gene for a GFP-like protein can be linked to genetic sequences for either a peptide or an entire protein. Upon

5 expression within a cell, such a chimeric molecule has the fluorescent properties of GFP along with the properties of the linked peptide or protein. When the linked peptide or protein is a substrate, the result is a protein-based fluorescent molecular substrate. Such a chimeric molecule can be produced in bacteria or cultured eukaryotic cells and purified for use as an exogenous

10 reporter substrate, or the gene can be introduced into the cell of interest for the endogenous production of reporter substrate. In fact, such a gene can be introduced into a cell-line to produce a new cell-line that carries the gene from one generation to the next. After intracellular turnover and sub-cellular sampling, electrophoresis can be used to separate the GFP-linked turned-

15 over substrate molecules from non-turned over GFP-linked substrate molecules. It is clear that genes for fluorescent proteins exist in other organisms, such as *Renilla*, that are not identical to the GFP from *Aequorea*, and it is expected that when these genes are cloned they will be similarly utilized.

20 Detection of molecular reporter molecules, obtained from within a cell, need not be achieved only by fluorescent methods. Fluorescent methods are generally utilized because of the extremely low limits of detection that they provide. In fact, fluorescence-based methods have been developed that can allow the detection of single molecules. As illustrated in Table 1, the

25 necessary detection limit to observe the amount of substrate found in a 1 pl (picoliter) mammalian cell at a physiologically normal concentration of 1 μM , is 10^{-18} mol or about 600,000 molecules. Alternative quantitative optical spectroscopic detection methods based on phosphorescence or chemiluminescence can also achieve this low limit of detection. Also some

30 electroanalytical methods such as amperometry or voltammetry are sensitive enough to be used to detect and quantify altered and unaltered reporter

molecules obtained from single cells. Sensitive detection and quantitation has also been achieved by coupling electron ion-spray mass spectrometry to capillary electrophoretic separation. Incorporation of heavy atoms into the reporter molecules, either from elements rare to biological systems or in the
5 form of unusual isotopes, could greatly increase the utility of such mass spectrometry.

Table 1.

	MAMMALIAN CELL (RBL)	<i>XENOPUS</i> OOCYTE	CONVENTIONAL <i>IN VITRO</i> ASSAY*
Volume Assayed	1 μ l (10^{-12} liters)	22 μ l (2.2×10^{-11} liters)	37.5 μ l (3.75×10^{-5} liters)
Concentration of Substrate	1 μ M (10^{-6} mol/l)	1 μ M (10^{-6} mol/l)	10 μ M (10^{-5} mol/l)
Total Amount Measured	10^{-18} mol (~600,000 molecules)	2.2×10^{-17} mol (~13,300,000 molecules)	3.75×10^{-10} mol (~226,000,000,000,000 molecules)

*Promega (Madison, WI) PKC Assay—normal instructions followed

Multiple reporter substrates, each reporting on a specific chemical
 10 reaction, are expected to be utilizable simultaneously within a given cell.

Finally, it is anticipated that the use of reporter molecules, as disclosed
 herein, will have utility in contexts that are not exclusively intracellular. This
 approach will be generally applicable to the measurement of chemical
 activities in circumstances where minute volumes, low concentrations, and
 15 miniscule quantities occur. An example of a non-cellular volume that could be
 assayed with such reporter molecules is the interior of a liposome or vesicle,
 a single-cell sized, or smaller, compartment that is typically enclosed by lipid
 molecules.

The invention and its various embodiments, now having been
 20 summarized, may be better understood by viewing the following drawings
 wherein like elements are referenced by like numerals.

Brief Description of the Drawings

Figs. 1A-H are schematic illustrations, in stages, showing the
 25 measurement of an intracellular kinase according to the invention. A legend is
 included with Fig. 1A to identify the molecular species depicted.

Fig. 1A is a diagrammatic depiction of a cell in a possible state prior to
 measurement wherein the cell contains inactive kinase molecules.

Fig. 1B a diagrammatic depiction of the cell of Fig. 1A after introduction of reporter substrate molecules.

Fig. 1C is a diagrammatic depiction of the cell of Fig. 1B during extracellular treatment with a stimulant that leads to activation of the
5 intracellular kinase.

Fig. 1D is a schematic representation of the reporter substrate phosphorylation as catalyzed by an active kinase molecule.

Fig. 1E is a diagrammatic depiction of the cell of Fig. 1C after some degree of reporter substrate has been phosphorylated.

10 Fig. 1F is the cell of Fig. 1E after disruption of its plasma membrane to obtain the contents for analysis in Fig. 1G.

Fig. 1G represents the separation of nonphosphorylated and phosphorylated substrate molecules by capillary electrophoresis (CE).

Fig. 1H shows a typical electrophoretogram in which a population of
15 nonphosphorylated reporter substrate molecules has been separated from a population of phosphorylated reporter substrate molecules.

Figs. 2A-C is the graph of three stacked electrophoretograms of reporter substrate molecule separations all made under identical electrophoretic conditions on the same day.

20 Fig. 2A is the graph of an electrophoretogram showing the separation of nonphosphorylated and *in vitro* phosphorylated reporter substrate molecules from a mixture in buffer solution, to be viewed as a reference for the electrophoretic mobilities of both species.

Fig. 2B is the graph of an electrophoretogram of the contents of a rat
25 basophilic leukemia (RBL) cell, a tumor cell, that was microinjected and in which protein kinase C (PKC) was not specifically activated (Fig. 3B shows a similar treatment).

Fig. 2C is the graph of an electrophoretogram showing the separation from the contents of an RBL cell in which protein kinase C (PKC) was
30 specifically activated.

Figs. 3A-C is the graph of three stacked electrophoretograms of

reporter substrate molecule separations all made under identical electrophoretic conditions on the same day.

Fig. 3A is the graph of an electrophoretogram showing the separation of nonphosphorylated and *in vitro* phosphorylated reporter substrate molecules from a mixture in buffer solution to establish a standard for the electrophoretic mobilities of both species.

Fig. 3B is the graph of an electrophoretogram showing the separation from the contents of a rat basophilic leukemia cell in which PKC was not specifically activated.

10 Fig. 3C is the graph of an electrophoretogram showing the separation from the contents of an RBL cell in which PKC was activated.

Figs. 4A-D is the graph of four stacked electrophoretograms of reporter substrate molecule separations all made under identical electrophoretic conditions on the same day. (The electrophoretic conditions differ from those used in Figs. 2 A-C and 3 A-C.)

Fig. 4A is the graph of an electrophoretogram that shows the mobility of the nonphosphorylated form of the reporter substrate in buffer.

Fig. 4B is the graph of an electrophoretogram that shows the mobility of the phosphorylated form of the reporter substrate in buffer.

20 Fig. 4C is the graph of an electrophoretogram that shows the separation of a sample of the contents of a *Xenopus* oocyte that had been previously microinjected to contain a reporter substrate.

Fig. 4D is the graph of an electrophoretogram that shows the separation of a sample of the contents of a *Xenopus* oocyte that had been previously microinjected to contain a reporter substrate which has been specifically activated.

Fig. 5 is an electrophoretogram showing the separation of a sample of the contents of a *Xenopus* oocyte that had been previously microinjected to contain a reporter substrate. The oocyte was treated to show an intermediate degree of activation.

Fig. 6A is a graph of the percentage of phosphorylation as a function of

time showing the intracellular phosphorylation of a reporter substrate in response to extracellular treatment with a potent pharmacologic specific activator.

Fig. 6B is a graph of phosphorylation as a function of time showing the time course of intracellular phosphorylation of reporter substrate in response to extracellular treatment of *Xenopus* oocytes with a physiological activator.

Fig. 7 is an electrophoretogram showing the separation of three different specific kinase reporter substrates from a sample of the contents of a *Xenopus* oocyte that had been previously microinjected.

10 Figs. 8A and 8B are diagrammatic depictions of the components of the apparatus used to make measurements of intracellular enzyme activity.

Fig. 8A is a diagram of a device for holding and sampling an oocyte (or similar cell) and rapidly subjecting the sample to electrophoresis.

Fig. 8B is a diagram of a microscope, capillary, and fluorescence
15 detection setup employing the device of Fig. 8A.

Fig. 9 is a diagrammatic depiction of the fast lysis apparatus used in the illustrated embodiment.

Fig. 10 is a diagrammatic depiction of the cell chamber, holding mammalian cells and electrophoresis buffer, used in conjunction with the fast
20 lysis apparatus in the illustrated embodiment.

Fig. 11A-C is the graph of three stacked electrophoretograms of reporter substrate molecule separations all made under identical electrophoretic conditions on the same day.

Fig. 11A is the graph of an electrophoretogram showing separation of
25 nonphosphorylated and *in vitro* phosphorylated reporter substrate molecules from a mixture in buffer solution as a reference for the electrophoretic mobilities of all species of reporter molecules.

Fig. 11B is the graph of an electrophoretogram of the contents of an RBL cell in which PKC was not specifically activated.

30 Fig. 11C is the graph of an electrophoretogram of the contents of an RBL cell in which PKC was activated.

Fig. 12 is a graph of the percentage of phosphorylation as a function of time showing the intracellular phosphorylation of a reporter substrate, in individual murine fibroblast NIH/3T3 cells, in response to extracellular treatment with a potent pharmacologically specific activator of PKC.

5 The invention and its various embodiments may now be understood by turning to the following Detailed Description of the Preferred Embodiments.

Detailed Description of the Preferred Embodiments

In the first illustrated embodiment the net activity of an intracellular enzyme known as protein kinase C (PKC) is measured by first introducing
10 into a single mammalian cell of ~1 pl volume, by means of microinjection, a synthetic fluorescent peptide that is a substrate for phosphorylation by PKC. Upon phosphorylation by PKC, the net charge of the fluorescent peptide substrate is altered from +5 to +3. More significantly, the charge-to-mass ratio of the peptide substrate changes as a result of this chemical alteration.

15 Consequently, when the contents of the single cell are obtained by means of fast, controllable laser lysis, separated by capillary electrophoresis, and detected by laser-induced fluorescence, one or two identifiable populations of molecules are observed and quantifiable. The populations are identified by reference to the characteristic migration time observed for each
20 molecular species when individually subjected to capillary electrophoresis. The faster migrating population of molecules, observed as a peak of induced fluorescence by a photomultiplier tube, corresponds to substrate that is not phosphorylated. The slower population, also observed as a peak of induced fluorescence, corresponds to substrate that has been phosphorylated. This
25 difference in migration rate through a capillary 22 is referred to as a change in electrophoretic mobility. The difference in electrophoretic mobility between the two populations of molecules is a consequence of the difference in the charge-to-mass ratio of the molecules in each population and the particular conditions used to carry out electrophoresis.

30 In this case, PKC phosphorylation of a subset of substrate molecules has altered their characteristic electrophoretic mobility. The relative amounts

of the molecules within the two populations give a measure of the net PKC activity that occurred within the single cell up to the time of lysis. Where the cell is not treated to activate its PKC, most of the detectable substrate peptide is in the nonphosphorylated form. When the cell is exposed to chemicals that
5 are activators of PKC for five minutes, approximately half the measured substrate peptide is of the phosphorylated form. These measurements have been taken from individual cells of two unrelated mammalian cell lines, rat basophilic leukemia (RBL) cells, a cultured granulocyte cell line, and NIH/3T3 cells, a mouse (murine) fibroblast cell line. Further in this embodiment, it is
10 demonstrated that two different fluorescent peptides that are reporters for the activity of two distinct kinases can be simultaneously detected and identified from the contents of a single RBL cell.

In another embodiment, the net activity of PKC within a single frog (*Xenopus*) oocyte **46'**, an approximately 1 mm in diameter, 1 μ l in volume,
15 nonmammalian cell, is measured. In this case, the same synthetic fluorescent peptide substrate is introduced by injection into the cell. Subsequently, a portion of the contents of the cell are removed with a sharpened capillary tip that is guided by a piezoelectric motor. In this case, the contents of the oocyte **46'** need not be harvested by laser lysis because the oocyte **46'** is large
20 enough to be mechanically sampled. Once a portion of the contents are in the capillary **22**, they are separated under electrophoretic conditions that have been empirically optimized for the separation of oocyte cytosolic components. Although differing cells and electrophoretic conditions are used as compared to the first embodiment, identifiable and quantifiable populations of
25 phosphorylated and nonphosphorylated PKC substrate molecules are still obtained. Additional measurements from the oocytes **46'** at multiple time points allowed the average rate of stimulated substrate phosphorylation to be determined for treatment of the oocytes **46'** with two different chemical activators of PKC.

30 Finally in this embodiment, it is demonstrated that three different fluorescent substrate peptides that are reporters for the activity of three

different distinct kinases can be detected and identified from a portion of the contents of a single frog oocyte 46'.

Shared Aspects of Two Illustrated Embodiments

Two preferred embodiments are presented. The first pertains to the measurement of intracellular chemical activity in single small mammalian cells that are typically 5-10 μm in diameter and about 1 pl in volume. This is illustrated by measurements made from individual rat basophilic leukemia (RBL) cells of the cell line RBL-2H3 and from individual murine NIH/3T3 cells. The second pertains to the measurement of intracellular chemical activity in relatively large single cells. In this second embodiment, measurements are made from individual *Xenopus laevis* frog oocytes 46' that are about 1 mm in diameter and about 1 μl in volume. In both illustrated embodiments, measurements of the activity of intracellular protein kinase C (PKC) are made from individual cells. In both embodiments, the same reporter substrate molecule is used, termed FI-sPKC for "fluorescent substrate of PKC". Upon phosphorylation by PKC, the net charge of the FI-sPKC is altered from +5 to +3. More significantly, the charge-to-mass ratio changes as a result of this chemical alteration. Subsequently, when the contents of the single cell are obtained, separated by capillary electrophoresis, and detected by laser-induced fluorescence, one or two identifiable populations of molecules, corresponding to nonphosphorylated and/or phosphorylated FI-sPKC, are observed and quantifiable. A comparison of the quantities involved (volume assayed, concentration of substrate, and total amount) in these PKC measurements and in a conventional radiographic method is provided in Table 1.

The sequence of amino acid residues that comprises this reporter peptide is derived from a region of the PKC protein sequence. The peptide sequence, as found in the larger PKC sequence, Arg-Phe-Ala-Arg-Lys-Gly-Ala-Leu-Arg-Gln-Lys-Asn-Val, is known to inhibit the enzyme, but when the underlined alanine residue is substituted with a serine residue, the resulting sequence is known to serve as a relatively specific substrate for PKC. Thus,

the sequence of FI-sPKC is FI- Arg-Phe-Ala-Arg-Lys-Gly-Ser-Leu-Arg-Gln-Lys-Asn-Val, where "FI" stands for a fluorescein moiety that was covalently attached to the amino terminal end of the peptide following standard peptide synthesis. The underlined serine residue, Ser, is the phosphorylation site.

5 Resin-bound, nonfluorescent sPKC peptide was obtained from the Stanford PAN facility and generated by conventional synthetic methods. The peptide with fluorescein covalently bound to the amino terminus was prepared by incubating 5-carboxyfluorescein succinimidyl ester (100-200 mg/ml, Molecular Probes, Eugene, OR) with resin-bound nonfluorescent peptide (30 mg/ml
10 peptide), 1-hydroxybenzotriazole (57 mg/ml), and N,N diisopropylethylamine (55 mg/ml) in dimethylformamide for 12 hours on a rotating mixer. The peptide was washed with dimethylformamide and ethyl acetate, and then cleaved from the resin by incubation for two hours with a trifluoroacetic acid (TFA)/free-radical scavenger mix (trifluoroacetic acid (84%, v/v), phenol
15 (0.2%, wt/v), thioanisole (5%, v/v), ethanediol (2.5%, v/v), water (8.5%, v/v)), previously sparged with argon or nitrogen. The peptide was separated from the resin by passage through a sintered glass filter. The peptide was then precipitated by addition of ice cold ether, dissolved in acetic acid (5%), and subsequently lyophilized. The peptide was purified by reverse phase high
20 pressure liquid chromatography (HPLC) on a semipreparative C18 column (Alltech, Deerfield, IL) with a gradient of 0.1% trifluoroacetic acid in water and 0.1% trifluoroacetic acid in acetonitrile. The concentration of the fluorescein-labeled peptide, FI-sPKC dissolved in buffer A (135 mM NaCl, 5 mM KCl, 10 mM HEPES, 2 mM MgCl₂, and 2 mM CaCl₂ adjusted to pH 7.4
25 with NaOH), was determined by comparing its fluorescence to that of standards of hydrolyzed carboxyfluorescein succinimidyl ester. FI-sPKC purity was assessed by reverse phase high pressure liquid chromatography on an analytic C18 column (Alltech) and by capillary electrophoresis. The peptide was stored in 2 µl aliquots at -70° C.

30 In both embodiments, reference separations, as illustrated in Figs. 2A, 3A, 4A and 4B, of the reporter substrate peptide, FI-sPKC, and its

phosphorylated form, referred to as P-FI-sPKC, were performed in buffer. Generation of P-FI-sPKC for this purpose was achieved by *in vitro* phosphorylation. FI-sPKC (4-40 μ M) was phosphorylated by rat brain PKC (1.2 μ g/ml, Molecular Probes, Eugene, OR) in the presence of

5 dioctanoylglycerol (25 μ g/ml, Avanti Polar Lipids, Alabaster, AL), phosphatidylserine (250 μ g/ml), $MgCl_2$ (10 mM), Tris HCl (50 mM, pH 7.5), adenosine triphosphate (1 mM), and Ca^{2+} (\sim 10 μ M). The free Ca^{2+} concentration was set by addition of 10 mM EGTA with 10 mM calcium. The reaction mixture was incubated at room temperature for one hour and the

10 phosphorylated peptide was used without further purification.

The general scheme of the measurements in both embodiments is depicted in Fig. 1A-H. Figs. 1A-H are schematic illustrations, in stages, showing the measurement of an intracellular kinase according to the invention. A legend is included with Fig. 1A to identify the molecular species

15 depicted. Fig. 1A is a diagrammatic depiction of a cell **46** in a possible state prior to measurement wherein the cell contains inactive kinase molecules **112**.

A Measurement

The course of a measurement begins with introduction of FI-sPKC into

20 the cell of interest, such as depicted in Fig. 1B. In this case, microinjection of a reporter substrate **114** through a micropipet **116** is depicted.

In measurements where the cell **46** is treated with a stimulant **118** to activate intracellular PKC, the stimulant **118** is added into a solution **120** surrounding the cell **46** as depicted in Fig. 1C.

25 Fig. 1D is a schematic representation of the reporter substrate phosphorylation as catalyzed by an active kinase molecule. Adenosine triphosphate (ATP) **124**, normally within the cell **46** at all times, is the intracellular source of phosphate groups for this reaction. The active kinase **122** catalyzes the reaction of the nonphosphorylated reporter substrate **114**

30 with ATP **124**. After phosphorylation, adenosine diphosphate (ADP) **126** is released along with the phosphorylated reporter substrate **128**.

Phosphorylation of FI-sPKC, as generally diagrammed in Fig. 1D, occurs to the greatest extent in cells that are treated with the stimulant **118**, resulting in cells **46** with internal contents as depicted in Fig. 1E. Fig. 1E is a diagrammatic depiction of the cell **46** of Fig. 1C after some amount of the reporter substrate has been phosphorylated. After a given time interval, in cases both with stimulant treatment and without, all or a portion of the intracellular contents of the cell **46** are loaded into a capillary **30** for separation and detection. Fig. 1F is a diagrammatic depiction of the cell **46** of Fig. 1E after disruption of its plasma membrane, shown in dotted outline, to liberate the cellular contents for analysis and for separation in the capillary **22** in Fig. 1G. Fig. 1G diagrammatically depicts the separation of nonphosphorylated and phosphorylated substrate molecules **114** and **128**, respectively, by capillary electrophoresis (CE) symbolically shown by the distance of separation between nonphosphorylated and phosphorylated substrate molecules **114** and **128**, respectively, within the capillary **30**.

An optical window **38** at the distal end of the capillary **30** allows interrogation by a laser (not shown) and collection of laser-induced fluorescence as represented in the combination Figs. 1F-G by flow of material from the cell **46** with a disrupted plasma membrane into the capillary **30**.

In both embodiments a photomultiplier tube (PMT) is used to collect argon laser (488 nm) induced fluorescence for detecting the presence and amount of FI-sPKC and P-FI-sPKC. Although the light collection systems used differ slightly, in both embodiments the photomultiplier tube current was amplified and converted to a voltage with a preamplifier, and this signal was digitized by a data acquisition board in a personal computer. The output is diagrammatically depicted in Fig. 1H, which shows a typical electrophoretogram in which a population of the nonphosphorylated reporter substrate molecules **114** has been separated from a population of the phosphorylated reporter substrate molecules **128**.

Fig. 9 is a diagrammatic depiction of the fast lysis apparatus used in the illustrated embodiment in which fast controlled cell lysis is performed in

combination with analysis by capillary electrophoresis. The system is described in the incorporated parent application, but the basic description is set forth again for convenience. It must be understood that many components now known or later devised may be substituted into the system of Fig. 1A to perform equivalent functions. For example, although the illustrated embodiment contemplates laser induced fluorescence detection of analytes in an electrophoretic column, any measurement device or detection mechanism capable of analyte detection or analysis may be used with any separation system capable of separating reporter molecules on the basis of charge, or charge-to-mass ratio.

The system, generally denoted by reference numeral **10**, is comprised of a microscope **12** having a video ocular readout **14** displayed on a CRT screen **16** or recorded by a videotape recorder or digital recorder (not shown). In particular, as will be described below, digital storage of the images and pattern processing in a computer system for automated cell processing and analysis is specifically contemplated as being within the scope of the present invention. A CCD (charge coupled device) video camera system **14, 16** recorded the real time bright field image of the cell **46** every 33 milliseconds. Lysis was determined from the appearance of the cell after the laser pulse. In most cases, the cell membrane was totally disrupted, leaving behind only a remnant of the membrane attached to a cover slip **36** with the cellular contents suspended in the immediate vicinity of the (former) cell. In the illustrated embodiment, electrophoresis was initiated by a voltage pulse sent by the laser at the time of operator-triggered firing and consequent lysis. It is to be expressly understood that computer-controlled pattern recognition could be substituted for operator viewing and firing could likewise be autonomously triggered.

A pulsed Nd:YAG laser **18** is directed through a microscope objective **20** of a microscope **12**. The laser system used for the cell lysing experiment included a frequency doubled Q-switched Nd:YAG laser such as manufactured as the Surelite I, by Continuum of Santa Clara, California. The

laser was used to generate a single laser pulse of 1 to 25 μJ with a 5 nsec pulse with a 532 nm wave length. The laser beam from the laser 18 was directed into the microscope 12, an Axiovert 135 manufactured by Zeiss of Thornwood, New York. The laser pulse was focused to approximately 0.3 to 0.4 μm at its waist 50 as shown in Fig. 1B using the microscope objective 20 (100 x, 1.3 n.a. Zeiss) at the interface of the cell chamber cover slip 36 and buffer solution 54. The cell 46 which was to be lysed was positioned 10-20 μm laterally to the focal point 50 of the laser pulse. Lysis, disruption of the cell's outer membrane, is effected by a shock wave that is generated by the laser pulse. It is to be expressly understood, however, that in another embodiment direct interaction may be arranged, such as where the laser parameters are adjusted only to open a hole in the cell membrane to expel the cytoplasmic contents, leaving the nuclear material within. It must be understood that other types of lasers other than the one illustrated here may also be used to generate the shock wave. In addition to lysis of the cell, the nucleus of the cell which has been isolated or removed from the cell may be manipulated to recover the DNA or other nuclear material.

A fused silica capillary column 22 is positioned by means of a micromanipulator 24 which positions a proximate inlet 26 of the capillary 22 above a cover slip or slide 28 positioned on the microscope stage 30. The buffer solution around the cell and above the cover slip is electrically grounded. A high voltage power supply, model CZE 1000R manufactured by Spellman of Plainview, New York, was used to drive the electrophoresis in the column or capillary 22. The fused silica capillary 22 had a 50 μm inner diameter and 360 μm outer diameter. The total length of the capillary 22 was 90 to 100 cm. A detection window 38 was about 75 cm from the inlet end 26. Electrophoresis was performed in a biologically compatible buffer. The cell chamber 199 of Fig. 10 served as an inlet reservoir and was held at ground potential. The outlet reservoir was held at 10 to 18 kV. The inlet end 26 of the capillary 22 was used as a micropipette for introducing the cellular contents into the capillary 22 after cell lysis.

After removing 5 mm of polyimide coating from the capillary **22** above the inlet end **26**, the inlet end **26** was mounted perpendicularly to the cover slip **36** on a micromanipulator **24**. The micromanipulator **24** enabled precise positioning of the capillary lumen **52** with respect to the cell **46** to be lysed and loaded into the capillary **22**. The capillary **22** was washed after every run. Gravitational siphon fluid flow was used to load the capillary **22** with phosphorylated/nonphosphorylated peptide standards. The loaded volumes were calculated from Poiseuille's equation.

The capillary **22** includes the optical detection window **38** upon which an excitation beam from the Argon laser **40** is aimed. Fluorescence data were collected at a right angle to the capillary **22** and to the laser beam from the laser **40** by a 40X microscope objective with a 0.75 n.a., Plan Fluor manufactured by Nikon of Melville, New York. The collected light was measured by a photomultiplier tube, a PMT R928 made by Hamamatsu, of Bridgewater, New Jersey, after spectral filtering through a 488 nm notch filter, manufactured by Kaiser Optical Systems of Ann Arbor, Michigan, and a band pass filter 535 DF55, made by Omega Optical of Brattleboro, Vermont. The photomultiplier current was amplified and converted to a voltage with a preamplifier then digitized by a data acquisition board in a personal computer **44**. The data were plotted and peak areas calculated using Origin written by Microcal of North Hampton, Massachusetts. Induced fluorescence data are used to create the electrophoretograms of Figs. 2A-C, 3A-C, 4A-D, 5, 7, and 11. The bath **34** is electrically coupled to a high voltage source **36** to provide the electrophoretic force to cause analyte separation in the capillary **22**.

*Conditions and Results Unique to the First Embodiment**The Cells Used*

Figs. 2A-C and 3A-C are two sets of graphs of three stacked electrophoretograms of reporter substrate molecule separations all made under identical electrophoretic conditions on the same day. In the first embodiment, illustrated by Figs. 2A-C and Figs. 3A-C, rat basophilic leukemia (RBL) cells, from the tumor mast cell line RBL-2H3, were used. These cells were grown at 37° C and 5% CO₂ in Dulbecco's modified eagle medium (DMEM) supplemented with 10% fetal bovine serum, and L-glutamine (584 g/l). Penicillin (100 units/ml) and streptomycin (100 µg/ml) were added to the media to inhibit bacterial growth. The cells 46 were grown in a chamber 199 made by using Sylgard (Dow Corning, Midland, MI) to attach a silicon "O" ring (15/16 in. outer diameter - not shown) to a 25 mm, round, no. 1, glass cover slip (not shown). Cell attachment to the glass surface was enhanced by coating the cover slip with Cell-Tak (Becton Dickinson, Bedford, MA) prior to adding the cells 46 to the cell chamber 199. Prior to use, the cells 46 were allowed to grow in supplemented media for 12-24 hr after plating in the cell chamber 199. The cells 46 were plated at concentrations determined empirically to produce approximately one cell per 100x field of view on the day of the measurement. Tissue culture materials were obtained from Gibco BRL (Gaithersburg, MD). Murine NIH/3T3 cells, as used in Fig. 12, were grown under identical conditions except that the use of Cell-Tak was omitted and the cells were typically allowed to grow 24-48 hours after plating in the cell chamber 199. Murine NIH/3T3 cells grow to approximately the same dimensions as do RBL cells.

Microscopy and Cell Sampling

In the first embodiment, the microscopy system of Fig. 9 used for cell lysis was similar to that described previously in the incorporated application entitled "Fast Controllable Laser Lysis of Cells for Analysis". The frequency-doubled Q-switched Nd:YAG laser **18** (Minilite II, Continuum, Santa Clara, CA; Minilase II-20, New Wave Research, Sunnyvale, CA) was used to generate a single pulse (1-25 μJ , 5-ns pulse width, 532 nm) directed into a fluorescence microscope **12** (Axiovert 135, Zeiss, Thornwood, NY). The pulse was focused ($\sim 0.3\text{-}0.4\ \mu\text{m}$ at its waist) with a microscope objective **20** (100x, 1.3 n.a., Fluor, Zeiss) within the cover slip **36** adjacent to its interface with the buffer. This same microscope objective **20** was used for visual observation during cell microinjection and selection. The cell **46** to be lysed was positioned 10 μm lateral to the focal point of the laser pulse so that no direct interaction of the laser beam with the cell **46** occurred. Lysis was effected by manual triggering of a Nd:YAG laser pulse. In most cases, the cell membrane was completely disrupted leaving behind only a remnant of the cell membrane attached to the cover slip.

Separation Conditions

In the first embodiment, electrophoresis was performed in a 50-75 cm long, fused-silica, uncoated capillary **22** (360 μm outer diameter, 50 μm internal diameter) from Polymicro Technologies (Phoenix, AZ) using buffer A, a physiologically compatible extracellular buffer composed of 135 mM NaCl, 5 mM KCl, 10 mM HEPES, 2 mM MgCl_2 , and 2 mM CaCl_2 and adjusted to pH 7.4 with NaOH. The cell chamber **199** served as the inlet reservoir and was held at ground potential. An outlet reservoir was held at a negative voltage of 10.8-13.4 kV with a high-voltage power supply (CZE100OR, Spellman, Plainview, NY). The electrophoretic current was 50-70 μA . The inlet **26** of the capillary **22** was used as a micropipet for introducing the cellular contents into the capillary **22** following cell lysis. Electrophoresis was initiated simultaneously with lysis by electrically triggering the voltage difference off

the laser pulse. The inlet and outlet ends **26** and **32** respectively of the capillary **22** were held at the same height during cell lysis, loading and electrophoresis to eliminate flow due to gravity. Between electrophoretic runs, the capillary **22** was washed by the following regimen:

- 5 1) buffer A was manually forced though the capillary **22** with
a handheld syringe for 30 sec (approximately 10 column volumes);
- 2) doubly distilled, reverse-osmosis treated water (Milli-Q)
was forced through for 30 seconds;
- 3) 1 N NaOH was forced through for 5 min.;
- 10 4) Milli-Q water was forced through for 30 sec.;
- 5) 0.1 N HCl was forced through for 3 min.;
- 6) Milli-Q water was forced though for 30 sec.; and
- 7) buffer A was forced though for 1 min.

Detection

For the purpose of detection, a 5 mm length of the polyimide coating was removed from the outer surface of the capillary 22 at a position 12.5 cm from the outlet end 32 to create a window 38. The coating was removed by
5 brief heating with a disposable butane lighter and was subsequently cleaned with a 70% ethanol solution. The capillary 22 lumen was interrogated through the optical window 38 by the focused beam of an argon ion laser (488 nm, Uniphase, San Jose, CA). The laser beam was focused by a lens (not shown) with a focal length of 3.8 cm (CVI, Albuquerque, NM). Fluorescence data
10 were collected by a detector 42 at a right angle to the capillary 22 and the laser beam with a microscope objective (not shown, but included in the detector 42 - 40x, 0.75 n.a., Plan Fluor, Nikon, Melville, NY), and the light measured with a photomultiplier tube (not shown, but included in the detector 42 - PMT, R928, Hammamatsu, Bridgewater, NJ) after spectral filtering with a
15 488 nm notch filter (Kaiser Optical Systems, Ann Arbor, MI) (not shown, but included in the detector 42) and a 535DF55 band-pass filter (Omega Optical, Brattleboro, VT) (not shown, but included in the detector 42). The photomultiplier tube current was amplified and converted to a voltage with a model 1212 preamplifier (DL Instruments, Dryden, NY) (not shown, but
20 included in the detector 42). The signal was digitized by a data acquisition board (DAS-1800, Keithly Metrabyte, Taunton, MA) in a personal computer 44 (Gateway, Sioux City, SD).

The data were plotted and peak areas calculated using Origin (Microcal, Northampton, MA). The system could easily detect $\sim 6 \times 10^{-21}$ mol
25 (about 3600 molecules) of FI-sPKC, well below the limit of detection required for physiologically relevant measurement (see Table 1).

Electrophoretic Migration Standards

Figs. 2A and 3A are two graphs of an electrophoretogram showing the separation of nonphosphorylated and *in vitro* phosphorylated reporter
30 substrate molecules from a mixture in buffer solution, to be viewed as a reference for the electrophoretic mobilities of both species. To obtain the

electrophoretic migration standards as shown in Figs. 2A and 3A, dilute mixtures of FI-sPKC and P-FI-sPKC were loaded into the capillary 22 by gravitational fluid flow and the loaded volume calculated from Poiseuille's equation. These mixtures were approximately 1 nM in concentration of both species in Fig 2A and approximately 100 pM in concentration of both species in Fig. 3A. Figs. 2A and 3A each show a nonphosphorylated FI-sPKC peak 136 and a phosphorylated P-FI-sPKC peak 138 for the standards.

Fig. 11A is also a graph of an electrophoretogram showing the separation of nonphosphorylated and *in vitro* phosphorylated reporter substrate molecules from a mixture in buffer solution. To obtain the electrophoretic migration standards as shown in Fig 11A, a dilute mixture of FI-sPKC, FI-sPKA, and P-FI-sPKC, each at approximately 1 nM in concentration, was loaded into capillary 22 by gravitational fluid flow and the loaded volume calculated from Poiseuille's equation. Fig 11A shows a nonphosphorylated FI-sPKC peak 136, a nonphosphorylated FI-sPKA peak 137, and a phosphorylated P-FI-sPKC peak 138 as standards.

Technical Aspects of Measurements in Figs. 2A - C

In the measurements illustrated by Fig. 2A - C, prior to mounting on the microscope stage for microinjection and analysis at room temperature (20°-23° C), each cell chamber 199 was treated by addition of 10 µl of 10 mM Pefabloc (Boehringer-Mannheim, Indianapolis, IN) in buffer A with gentle manually pipetted mixing using a Pipetman® (Rainin, Woburn, MA).

This yielded a final Pefabloc concentration of 200 µM. Pefabloc is a soluble, physiologically tolerated, inhibitor of serine proteases, a subset of intracellular enzymes that tend to degrade peptides. Pefabloc was included to minimize intracellular degradation of FI-sPKC. In instances where PKC was to be activated, as in Fig. 2C, 10 µl of 5 µM phorbol myristic acid (PMA, purchased from Sigma, St. Louis, MO.) freshly diluted from a 200 µM stock solution, into buffer A, was added and mixed gently at the same time as Pefabloc. The stock solution was composed of PMA dissolved in dimethyl sulfoxide (DMSO). The cell chambers 199 were returned to the incubator (37°

C, 5% CO₂), for 5 minutes. Immediately after mounting on the microscope stage **30**, the culture medium in the cell chamber **199** was replaced by continuous flow and removal (2 ml/min, approximately four cell chamber volumes/min) of room temperature (20°-23° C) buffer A with added 10 mM glucose and 200 µM Pefabloc. Flow was maintained throughout microinjection and recovery. This typically lasted 20-30 minutes. Flow was halted briefly during lysis, restarted 1 minute after lysis and capillary loading and continued throughout the course of electrophoresis.

In the measurements illustrated by Figs. 2A-C, FI-sPKC was diluted in buffer B, a physiologically compatible intracellular buffer composed of 10 mM Hepes and 140 mM KCl, at pH 7.4. The final concentration of FI-sPKC was ~800 µM. Mixed with the FI-sPKC was Calcium Orange (Molecular Probes, Eugene, OR) at a concentration of 500 µM. This mixture was microinjected into several cells in the same cell chamber through a Femtotip I glass microinjection needle **116** with an Injectman 5171 micromanipulator and a Transjector 5346 system at a pressure of 80 hPa for 0.2 sec (Femtotip, Injectman and Transjector are products of Eppendorf, Hamburg, Germany). The intracellular concentrations are estimated to be 8 µM FI-sPKC and 5 µM Calcium Orange. The injection was performed on the same microscope **12** as used for cell selection and analysis, namely an Axiovert 135 (Zeiss). The cells **46** that had been successfully microinjected were identified by visual observation of internal Calcium Orange epi-fluorescence through a set of TRITC filters (Chroma Technology Corp., Brattleboro, VT) (not shown, but included in the microscope **12**) to minimize photodestruction of the fluorescein moiety on FI-sPKC. Only the cells **46** that appeared healthy as judged by the presence of a uniformly fluorescent, homogeneous cytoplasm and smooth plasma membrane, were chosen for analysis. A large fraction of microinjected cells failed this requirement, demonstrating membrane defects, punctuate fluorescence, or excessively bright nuclei.

30 Results of Measurements in Figs. 2A - C

Fig. 2B is the graph of an electrophoretogram showing the separation

from the contents of a rat basophilic leukemia (RBL) cell **46** in which the enzyme protein kinase C (PKC) was not specifically activated (peak **144** at ~1200 sec. is thought to be a byproduct of intracellular proteolysis). In particular Fig. 2B is an electrophoretogram of the contents of an RBL cell **46** that was microinjected, as described above, in the absence of an extracellular stimulant. Fig. 2B shows a sizable nonphosphorylated fluorescent peak **140** that corresponds in migration time to FI-sPKC peak **136** shown in Fig. 2A. In contrast, only a very small peak **142** that corresponds in migration time to phosphorylated P-FI-sPKC is observed. Virtually no net intracellular PKC activity occurred within the treated cell **46** from the time of microinjection.

Fig. 2C is the graph of an electrophoretogram showing the separation from the contents of an RBL cell **46** in which PKC was specifically activated by treatment with 100 nM phorbol myristic acid (PMA). In contrast to the results of Fig. 2B, the electrophoretogram in Fig. 2C of the contents of an RBL cell **46** exposed to the extracellular, membrane permeant stimulant, PMA shows evidence of significant net PKC activity. The peak **146** corresponding to FI-sPKC and the peak **148** corresponding to P-FI-sPKC are of nearly equal magnitude.

Clearly activation of an intracellular kinase by a potent pharmacologic activator can be measured by this approach. Furthermore, both electrophoretograms in Figs. 2B and 2C show the presence of peaks that cannot be readily identified. It is believed that these peaks correspond to forms of FI-sPKC that have been partially degraded by intracellular proteases or covalently modified by other intracellular nonkinase enzymes. The occurrence of these molecular products as discrete, well defined peaks suggests that they represent discrete molecular species and are the products of a limited number of specific side reactions. The FI-sPKC molecules are not overwhelmingly consumed by these side reactions, since in both Figs. 2B and 2C, side products account for only about 50% of collected fluorescence. The unique species that migrates most slowly in Fig. 2C indicated as a peak **152** may correspond to a modified version of P-FI-sPKC.

Technical Aspects of Measurements in Figs. 3A - C

Fig. 3A is the graph of an electrophoretogram showing the separation of nonphosphorylated and *in vitro* phosphorylated reporter substrate molecules from a mixture in buffer solution, to be viewed as a reference for the electrophoretic mobilities of both species. Fig. 3B is the graph of an electrophoretogram showing the separation from the contents of a rat basophilic leukemia (RBL) cell **46** in which the enzyme protein kinase C (PKC) was not specifically activated. Fig. 3C is the graph of an electrophoretogram showing the separation from the contents of an RBL cell **46** in which PKC was activated as a consequence of treatment of the cell with extracellular ATP.

In the measurements illustrated by Figs. 3A-C the sequence of events is as follows. First, the cell chamber **199** was removed from an incubator, the culture medium replaced with two manually pipetted exchanges of buffer A containing additional 10 mM glucose, and the cells **46** within the chamber **199** microinjected, as described above, with a mixture of 100 μ M FI-sPKC and 500 μ M rhod-2 (Molecular Probes, Eugene, OR) in buffer B. The intracellular concentrations are estimated to have been 1 μ M FI-sPKC and 5 μ M rhod-2. The chamber **199** was then placed back in the incubator (37° C, 5% CO₂) for 15 minutes. After this recovery period all further manipulations occurred at room temperature (20°-23° C). The buffer within the cell chamber **199** was replaced with five exchanges of manually pipetted buffer A containing additional 10 mM glucose and placed back on the microscope stage for cell selection and analysis. The cells **46** that had been successfully microinjected were identified by visual observation of internal rhodamine epi-fluorescence through a set of TRITC filters (Chroma) in order to minimize photodestruction of the fluorescein moiety on FI-sPKC. Only those cells of the cells **46** that appeared healthy as judged by the presence of a uniformly fluorescent, homogeneous cytoplasm and smooth plasma membrane, were chosen for analysis. As described previously, a large fraction of the microinjected cells failed this requirement, demonstrating membrane defects, punctuate

fluorescence, or excessively bright nuclei.

In instances where PKC was to be activated, as in Fig. 3C, 10 µl doses of 500 µM adenosine triphosphate (ATP, Sigma, St. Louis, MO.) in buffer A were pipetted manually directly above the cell of interest. RBL-2H3 cells 46
5 express transmembrane purinergic receptors on their surface. When these receptors bind ATP, an intracellular G-protein linked signal transduction cascade becomes activated that can give rise to intracellular PKC activation. Finally, after cell selection and either no further treatment, as in Fig. 3B, or treatment with ATP, as in Fig. 3C, lysis was triggered by manual firing the
10 Nd:YAG laser 18 and electrophoresis started simultaneously as described above.

Results of Measurements in Figs. 3A - C

Fig. 3B is an electrophoretogram of the contents of an RBL cell 46 that was microinjected, as described above, and received no treatment with ATP.
15 The distribution of fluorescent peaks is very similar to that seen in Fig. 2B. The peak 154 that corresponds to FI-sPKC is six times greater in area than the peak 156 that corresponds to P-FI-sPKC. This indicates that six-fold more of the reporter substrate was nonphosphorylated as compared to phosphorylated. In the absence of treatment to activate PKC, as in Fig. 2B,
20 the net activity of PKC is low.

Fig. 3C is an electrophoretogram of the contents of an RBL cell 46 that was similarly microinjected, but subsequently exposed to two doses of ATP. The doses were administered 3 minutes apart with the last dose being added about 2 minutes prior to lysis. Fig. 3C demonstrates that a nearly 1:1 ratio of
25 FI-sPKC integrated under the peak 158 and P-FI-sPKC integrated under the peak 160 occurred within the cell 46 following extracellular treatment with ATP, a physiological activator of PKC.

Technical Aspects of Measurements in Figs. 11A - C

Fig. 11A is the graph of an electrophoretogram showing the separation
30 of nonphosphorylated and *in vitro* phosphorylated reporter substrate

molecules from mixture in buffer solution, to be viewed as a reference for the electrophoretic mobilities of each species, FI-sPKC, FI-sPKA, and P-FI-sPKC. At the time of this analysis, no *in vitro* phosphorylated P-FI-sPKA was available for use as a migration standard. Fig. 11B is the graph of an
5 electrophoretogram showing the separation from the contents of an RBL cell 46 in which the enzymes PKC and protein kinase A (PKA) were not specifically activated. Fig. 11C is the graph of an RBL cell 46 in which PKC was activated as a consequence of treatment of the cell with extracellular PMA.

10 In the measurements illustrated by Figs. 11B-C, the following compounds were mixed in buffer B: FL-sPKC at 120 μ M, FI-sPKA at approximately 100 μ M, and Calcium Orange (Molecular Probes, Eugene, OR) at 650 μ M. The mixture was microinjected at room temperature (20°-23° C) into several cells in the same cell chamber through a Femtotip I glass
15 microinjection needle 116 (Eppendorf, Hamburg, Germany) with a Micromanipulator 5171 and Transjector 5246 system (Eppendorf) at a pressure of 80 hPa for 0.2 sec. The intracellular concentrations are estimated to be 1.2 μ M FI-sPKC, 1 μ M FI-sPKA, and 6.5 μ M Calcium Orange. The injection was performed on the same microscope 12 as used for cell selection
20 and analysis, namely an Axiovert 135 (Zeiss). The cells 46 that had been successfully microinjected were identified by visual observation of internal Calcium Orange epi-fluorescence through a set of TRITC filters (Chroma Technology Corp., Brattleboro, VT) (not shown, but included in the microscope) to minimize photodestruction of the fluorescein moiety on the
25 reporter substrate molecules. Only the cells that appeared healthy as judged by the presence of a uniformly fluorescent, homogeneous cytoplasm and smooth plasma membrane, were chosen for analysis. A large fraction of microinjected cells failed this requirement, demonstrating membrane defects, punctate fluorescence, or excessively bright nuclei.

30 In cases where the cells were to be stimulated with PMA as in Fig. 11C, prior to injection, the culture media in the cell chamber 199 was replaced

by a solution of 1 μ M PMA in buffer A containing 10 mM glucose. The cell chamber was returned to the incubator (37° C, 5% CO₂) for 10 minutes.

In all cases, immediately after removal from the incubator, the cell chambers **199** were mounted on the microscope stage **30** and the solution in the cell chamber was replaced by continuous flow and removal (2 ml/min, approximately four cell chamber volumes/min) of room temperature buffer A with added 10 mM glucose. Flow was maintained throughout microinjection and recovery. This typically lasted 5-15 minutes. Flow was halted briefly during lysis, restarted within 1 minute after lysis and capillary loading and continued throughout the course of electrophoresis.

Results of Measurements in Figs. 11A - C

Fig. 11B is the graph of an electrophoretogram showing the separation from the contents of an RBL cell **46** in which the enzymes PKC and PKA were not specifically activated. In particular Fig. 11B is an electrophoretogram of the contents of an RBL cell that was microinjected, as described above, in the absence of extracellular PMA. Fig. 11B shows sizable peaks that correspond in migration times to FI-sPKC and FI-sPKA. No peaks that could be positively identified as the phosphorylated forms, P-FI-sPKC or P-FI-sPKA, of the reporter substrate peptides occurred in this trace. No measurable net intracellular PKC or PKA activity occurred within the cell **46** from the time of microinjection.

Fig. 11C is the graph of an electrophoretogram showing the separation from the contents of an RBL cell **46** in which the enzyme PKC was specifically activated with PMA. PMA is well characterized as a specific activator of PKC but not of PKA. As described above, this cell was microinjected after treatment with PMA. Fig. 11C shows sizable peaks that correspond in migration times to FI-sPKC, FI-sPKA, and a cluster of smaller peaks centered with a migration time that matches the phosphorylated form of FI-sPKC, P-FI-sPKC. All three chemical species separated from a mixture in buffer in Fig. **11A** are shown separated from the contents of a small mammalian cell in Fig. **11C**. Since a migration standard for the phosphorylated form of FI-sPKA was

not available at the time of this measurement, it is uncertain if any of these peaks correspond to P-FI-sPKA. As noted above, in the absence of stimulation with PMA, none of the peaks in the cluster centered around 910 seconds are seen at comparable magnitude. A small, but measurable amount of PKC activation has occurred within the cell analyzed in Fig. 11C.

Technical aspects of the Measurements in Fig. 12

Fig. 12 is a graph of degree of phosphorylation as a function of time showing the course of intracellular phosphorylation of reporter substrate by PKC in response to extracellular treatment of murine NIH/3T3 cells with 100 nM PMA, a potent and specific pharmacological activator of PKC. Each point represents the average of three or more measurements, and the error bars represent a single standard deviation.

In the measurements illustrated in Fig. 12 FL-sPKC at 300 μ M and Calcium Orange (Molecular Probes, Eugene, OR) at 900 μ M were mixed in buffer B. The mixture was microinjected at room temperature (20°-23° C) into several cells in the same cell chamber through a Femtotip I glass microinjection needle (Eppendorf, Hamburg, Germany) with a Micromanipulator 5171 and Transjector 5246 system (Eppendorf) at a pressure of 80 hPa for 0.2 sec. The final intracellular concentrations are estimated to have been 3 μ M for FI-sPKC and 9 μ M for Calcium Orange. Following microinjection, the cells were allowed to recover at room temperature for 10-15 minutes on the microscope stage with continuous flow (2 ml/min) of buffer A with added 10 mM glucose.

In cases where PKC was to be stimulated, PMA was diluted from a stock solution in dimethyl sulfoxide (DMSO) into buffer A containing additional 10 mM glucose immediately prior to use. This solution (approximately 500 μ l) was exchanged for the solution within a cell chamber manually with a hand-held Pipetman® (Rainin Instrument Co., Woburn, MA). This was carried out while the cell chamber was mounted on the microscope stage after the flow of exchange buffer had been paused. The flow of buffer was halted throughout the time of incubation with PMA containing solution. During

the time of PMA stimulation, a healthy cell was selected by observation of Calcium Orange epi-fluorescence. At the appropriate time point, either after 2.5 or 5 minutes, the cell was lysed by remote triggering of the Nd:YAG laser 18 and loaded into the inlet 26 of the capillary 22 by the simultaneous start of 5 electrophoresis.

The peak areas corresponding to FI-sPKC and P-FI-sPKC were measured using the software analysis package Origin 5.0 (Microcal Software, Inc., Northampton, MA).

Results of Measurements in Fig. 12

10 Fig 12 is a graph of degree of phosphorylation as a function of time showing the time course of intracellular phosphorylation of reporter substrate by PKC in response to extracellular treatment of murine NIH/3T3 cells with PMA, a potent and specific activator of PKC. Fig. 12 shows the quantitative results of a number of measurements conducted to determine the time course 15 of intracellular reporter substrate phosphorylation in NIH/3T3 cells in response to activation by exposure to 100 nM PMA at room temperature (20°-23° C). From the approximate correspondence of the 5 minute time point to the 50% phosphorylation level, a rate of FI-sPKC phosphorylation of 6 +/- 2 nM/s is estimated to occur within NIH/3T3 cells under these conditions. 20 At this rate, only 0.007% of the reporter substrate peptide would have been phosphorylated in the conservatively estimated 33 msec interval between laser induced cell lysis and the initiation of electrophoretic separation. Thus, the rapidity with which the contents of a cell are obtained and separated allows for exceptionally accurate snapshots of intracellular chemical reactions 25 to be obtained. Fig. 12 further demonstrates that measurements can be made from a second mammalian cell type, namely NIH/3T3 cells, as well as from RBL cells.

Conditions and Results Unique to the Second Embodiment

Summary

In the second embodiment, illustrated by Figs. 4A-8B *Xenopus laevis* frog oocytes **46'** were used as cells of interest. In the measurements of PKC activation, illustrated by Figs. 4A-6B, FI-sPKC was microinjected into an oocyte **46'**. Cytoplasm (~22 pl) from the oocyte **46'** was then loaded into the capillary **22** and electrophoresed. FI-sPKC and P-FI-sPKC were separated and quantified from their fluorescence. The ratio of P-FI-sPKC to FI-sPKC measures the relative activation of PKC, *i.e.* the intracellular balance between PKC and the phosphatases. Phosphatases catalyze the reverse of the reaction catalyzed by kinases—namely the removal of a phosphate from a substrate molecule.

Fig. 7 is a graph of phosphorylation as a function of time showing the separation of three different specific kinase reporter substrates from a sample of the contents of a *Xenopus laevis* oocyte **46'** that had been previously microinjected to contain ~1 μ M FI-sPKC, ~330 nM FI-sPKA, and ~10 nM FI-scdc2K. FI-sPKA, a specific reporter for protein kinase A (PKA) activity has the sequence FI-Lys-Arg-Arg-Glu-Ile-Leu-Ser-Arg-Arg-Pro-Ser-Tyr-Arg, and was derived from the CREB protein. FI-scdc2K, a specific reporter for cdc2 kinase (originally identified genetically as cell division cycle mutant 2) has the sequence FI-Gly-Gly-Gly-Arg-Ser-Pro-Gly-Arg-Arg-Arg-Lys, and comprises a consensus phosphorylation site derived from several proteins. The underlined serine residues are the sites of phosphorylation. The peptides were synthesized and labeled with fluorescein as described for FI-sPKC, except that FI-scdc2K was labeled with the mixed 5- and 6-isomers of carboxyfluorescein succinimidyl ester (100-200 mg/ml, Molecular Probes, Eugene, OR); thus, FI-scdc2k consisted of two isomeric forms. A peak **162** and a peak **172** were identified by their migration times as observed when injected into oocytes **46'** singly (not shown). The first doublet, peaks **162** and **164**, corresponds to two isomers of either phosphorylated or nonphosphorylated FI-scdc2K. The second doublet, peaks **166** and **168**,

corresponds to two isomers of the other form of FI-scdc2K. One peak 170 represents nonphosphorylated FI-sPKC, and one peak 172 represents nonphosphorylated FI-sPKA.

As shown in Fig. 7, when a plurality or mixture of peptides of different
5 kinase specificities was loaded into the oocyte 46', each of the different peptides could be detected. This demonstrates that from a single oocyte 46', it is possible to quantify simultaneously the activation of multiple kinases within a single pathway or in different pathways, making this technique a powerful tool to study cellular signal transduction cascades.

10 *Cells Used*

Oocytes 46' were surgically obtained from pigmented or albino *Xenopus laevis* frogs and were cultured as described previously. Oocytes 46' were microinjected by means of an injector mounted on a low power dissecting microscope (not shown) with 50 nl of reporter substrate peptide
15 (0.5 – 200 μ M) in buffer B, and then incubated for 30 min to permit diffusion of the peptide through the oocyte 46'. For measurements, oocytes 46' were placed in buffer C (96 mM NaCl, 2 mM KCl, 1.8 mM CaCl₂, 1 mM MgCl₂, 5 mM Hepes, pH 7.6). When LPA (1-oleoyl-2-hydroxy-sn-glycero-3-phosphate, Avanti Polar Lipids, Alabaster, AL) was added to the oocytes 46', fatty
20 acid-free bovine serum albumen (5%) was used as a carrier as described previously.

Separation Conditions

All samples from the oocytes 46' were obtained and electrophoresed
25 by a method referred to as piezoelectric sampling/rapid translation/capillary electrophoresis (PERT CE). The apparatus used for PERT CE is illustrated in Figs. 8A - B. Figs. 8A and 8B are diagrammatic depictions of the components of the apparatus used to make measurements of intracellular enzyme activity from single *Xenopus laevis* oocytes 46'. Fig. 8A shows a custom designed
30 holder 130 for the oocyte 46' and an electrophoresis buffer. The holder is part of a complete apparatus for automatically sampling oocyte cytoplasm and

rapidly subjecting this sample to electrophoresis before any substantial change in the substrate molecules can occur. This is achieved by providing a sharpened electrophoresis capillary **22** attached to a high speed translation system to plunge the capillary into the cell to capture a "plug of cytoplasm.

- 5 Then the capillary is automatically withdrawn from the oocyte and rapidly moved to make electrical contact with a reservoir **134** of electrophoresis buffer so that electrophoresis of the substrate molecules within the "plug" commences before any additional chemical reactions occur. In the preferred device a piezoelectric element moves the capillary up and down (vertical axis)
- 10 to sample the cell while a linear motor rapidly moves the capillary in a horizontal direction to transition from the oocyte to the reservoir **132**.

In Fig. 8A the arrows show the axes of movement of the capillary **22** by the piezoelectric element **P** and linear motor **LM**. A reservoir **132** with the oocyte **46'** contains a biologically compatible buffer C (96 mM NaCl, 2 mM

15 KCl, 1.8 mM CaCl_2 , 1 mM MgCl_2 , 5 mM Hepes, pH 7.6), while the second reservoir **134** contains the electrophoresis buffer. Fig. 8B shows a schematic of microscope, capillary, and fluorescence detection similar to Fig. 9. The holder **130** of Fig. 8A was mounted on the stage **174** of an upright microscope **176** or below a dissecting microscope. For simplicity, the

20 piezoelectric element and linear motor are not shown. The oocyte **46'** is placed in a "V"-shaped well in the reservoir **132** that had been drilled into a block of Teflon or polycarbonate **180**. The reservoir **132** was filled with the biologically compatible buffer, buffer C. A tip **182** of the separation capillary **22** (50 μm I.D., 360 μm O.D.) was etched to an outer diameter of $\sim 70\ \mu\text{m}$ by

25 treatment with hydrofluoric acid. The capillary tip **182** was mounted on the edge of a piezoelectric motor element (Piezo Systems, Cambridge, MA not shown) to allow controlled vertical movement of the tip **182** indicated by direction **P** in Fig. 8A. A micromanipulator (not shown) which held the piezoelectric device and the capillary **22** was mounted on a linear motor (not

30 shown) to allow controlled horizontal movement indicated by direction **LM** in Fig. 8A. Using the micromanipulator, the etched tip of the capillary **22** was

placed above and adjacent to the oocyte's plasma membrane.

To obtain cytoplasm, the capillary **22** was driven 60 μm into the oocyte **46'** by the piezoelectric element. This resulted in the filling of the tip **182** of the capillary **22** with a "plug" of sampled cytoplasm. The capillary **22** was then
5 automatically withdrawn from the cytoplasm. Immediately after withdrawing of the capillary **22** from the oocyte **46'**, the capillary **22** was moved horizontally (1.2 cm at a velocity of 6 cm/s) by the linear motor and then rapidly lowered by the piezoelectric element into a second reservoir **134** that contained an electrophoresis buffer. Electrophoresis was initiated immediately. This entire
10 sequence was computer controlled and accomplished in 350 ms. The linear motor (Oriental Motor Co., Torrance, CA) was controlled by a solid state relay board and an analog output board (Keithley, Cleveland, OH) (not shown).

Electrophoretic separations from oocyte cytoplasm were carried out under two distinct sets of conditions. In the measurements illustrated in Figs.
15 4A - D, capillaries with a proprietary neutral coating (Supelco, Bellefonte, PA) were used. In the measurements illustrated in Figs. 5 and 6A - B, polyethyleneimine (PEI) coated capillaries were used and prepared as follows. Fused silica capillaries **22** (Polymicro Technologies, Phoenix, AZ) were washed continuously for 30 min with 1 M NaOH or KOH followed by a
20 15-min wash with water. The capillaries **22** were then coated either by flowing PEI (5%) continuously through each capillary **22** for 60 min. or by loading each capillary **22** with PEI (5%) and then incubating the PEI-loaded capillary for 60 min. The solution of PEI was expelled from the capillary **22** with nitrogen. Each capillary **22** was rinsed with water for 15 min and then with the
25 electrophoresis buffer for 15 min.

All electrophoretic separations shown in Figs. 4A-7 are best understood as examples of the use of micellar electrokinetic chromatography (MEKC). This terminology refers to the use of an electrophoretic buffer in which detergent micelles are used to facilitate reproducible and efficient
30 separations. The use of micelles in these separations has proven necessary because *Xenopus* oocyte cytoplasm contains relatively large amounts of

amphiphilic molecules, particularly in the form of the highly negatively charged protein phosvitin. Electrophoresis was performed in a 50-100 cm long capillary **22** (360 μm outer diameter, 50 μm internal diameter). The electrophoretic voltage was 10-15 kV (current, 30-60 pA). The electrophoresis
5 was performed in buffer D (45 mM NaCl, 1.6 mM KCl, 0.6 mM MgCl_2 , 1% triton-X 100, and 10 mM Hepes (pH 7.4)), or buffer E (buffer D with 645 mg/ml cholic acid). Between electrophoretic runs, the capillary **22** was washed for 25 min with the electrophoresis buffer. The fluorescence of analyte bands migrating through the capillary **22** was measured 15-25 cm
10 from the outlet end of the capillary **22** as previously described. Peptide standards were loaded into the capillary **22** by gravitational fluid flow and the loaded volume calculated from Poiseuille's equation.

Detection

For detection purposes, a 5 mm length of the polyimide coating was
15 removed from the outer surface of the capillary **22** at a position 50 cm from the inlet tip **182**. The coating was removed by brief heating with a disposable butane lighter and was subsequently cleaned with a 70% ethanol solution. Through this optical window **38**, the capillary lumen was interrogated by the focused beam of an argon ion laser (operated at 488 nm) (Uniphase, San
20 Jose, CA). Fluorescence was collected at a right angle to the capillary **22** and laser beam by a collecting lens (not shown), and the light was measured with a R928 photomultiplier tube (PMT) (Hamamatsu, Bridgewater, NJ) (not shown) with a 533DF56 spectral filter (Omega Optical, Brattleboro, VT) (not shown). The photomultiplier tube current was amplified and converted to a voltage
25 with a preamplifier (not shown, but included in the detector **42**). The signal was digitized by a data acquisition board (DAS-1802ST-DA or DAS-1802HR-DA, Keithly Metrabyte, Taunton, MA) in a personal computer
44 (Gateway, Sioux City, SD). The data were plotted and peak areas calculated using Origin (Microcal, Northampton, MA). The detection limit of
30 this configuration was approximately 10^{-19} moles, about 60,000 molecules, well below the detection limit for physiologically relevant measurement (see

Table 1).

Technical Aspects of the Measurements in Figs. 4A - D

Fig. 4A is the graph of an electrophoretogram that shows the mobility of the nonphosphorylated form of the reporter substrate in buffer at a peak
5 184. Fig. 4B is the graph of an electrophoretogram that shows the mobility of the phosphorylated form of the reporter substrate in buffer at a peak 186. In Figs. 4A and 4B, reporter substrate standards were loaded into the capillary 22 by gravitational fluid flow and the loaded volume calculated from Poiseuille's equation. In the measurements illustrated in Figs. 4C and 4D,
10 individual oocytes 46' were microinjected with 50 nl of 200 μ M FI-sPKC as described and incubated in buffer C for 30 minutes to allow diffusion of the reporter substrate. The estimated final intracellular concentration of reporter substrate was \sim 10 μ M. Separations were achieved with PERT CE sampling followed by MEKC with buffer E through a capillary with a proprietary neutral
15 inner coating (Supelco, Bellefonte, PA). In the measurements, the reservoir 132 that contained the oocyte 46' was filled with buffer C, while the capillary 22, the second reservoir 134 that served as the inlet for electrophoresis and the output reservoir 34 contained buffer E.

Results of Measurements in Fig. 4A - D

20 Fig. 4C is the graph of an electrophoretogram that shows the separation of a sample of the contents of a *Xenopus laevis* oocyte that had been previously microinjected to contain \sim 10 μ M reporter substrate. In this case, the oocyte 46' was not treated to activate intracellular PKC. Fig. 4C is an electrophoretogram of a sample of cytoplasm from the oocyte 46' that was
25 not treated with an extracellular stimulant. Essentially all collected fluorescence at the peak 188 corresponded to nonphosphorylated FI-sPKC, as can be seen by reference to the peak 184 of Fig. 4A, which is an electrophoretogram of FI-sPKC loaded from buffer.

Fig. 4D is the graph of an electrophoretogram that shows the
30 separation of a sample of the contents of a *Xenopus laevis* oocyte that had

been previously microinjected to contain ~10 μ M reporter substrate. In this case, the oocyte **46'** was treated extracellularly with 10 μ M PMA for 10 minutes prior to sampling to specifically activate intracellular PKC. Fig. 4D is an electrophoretogram of a sample from the oocyte **46'** that was exposed to 10 μ M PMA for 10 minutes. In this case, essentially all collected fluorescence at the peak **190** corresponded to P-FI-sPKC as can be seen by reference to the peak **186** of Fig. 4B, an electrophoretogram of P-FI-sPKC loaded from buffer.

These results demonstrate that net activation of PKC in response to the potent pharmacologic stimulant PMA can be measured from a single *Xenopus* oocyte. Further, this shows that in the absence of stimulation, net PKC activity is relatively low in oocytes.

Technical Aspects of the Measurements in Figs. 5 and 6A, 6B

Fig. 5 is a graph of fluorescence as a function of time showing the separation of a sample of the contents of a *Xenopus laevis* oocyte **46'** that had been previously microinjected to contain ~10 μ M reporter substrate. In this case, the oocyte **46'** was treated with 10 μ M PMA for 3 minutes prior to sampling to specifically activate intracellular PKC. Both the nonphosphorylated and phosphorylated forms of the reporter substrate are present, indicating that substrate turnover was incomplete after three minutes.

Fig. 6A is a graph of phosphorylation as a function of time showing the time course of intracellular phosphorylation of reporter substrate by PKC in response to extracellular treatment of *Xenopus laevis* oocytes with 10 μ M PMA, a potent pharmacologic specific activator of PKC. Each point represents the average of three or more measurements, and the error bars represent a single standard deviation.

Fig. 6B is a graph of phosphorylation as a function of time showing the time course of intracellular phosphorylation of reporter substrate by PKC in response to extracellular treatment of *Xenopus laevis* oocytes with 10 μ M LPA (1-oleoyl-2-hydroxy-*sn*-glycero-3-phosphate, Avanti Polar Lipids, Alabaster, AL) a physiological activator of PKC. Each point represents the

average of three or more measurements, and the error bars represent a single standard deviation.

In the measurements illustrated in Figs. 5-7, individual oocytes **46'** were microinjected with 50 nl of 200 μ M FI-sPKC as described and incubated in buffer C for 30 minutes to allow diffusion of the reporter substrate. The estimated final intracellular concentration of reporter substrate was approximately 10 μ M. Separations were achieved with PERT CE sampling followed by MEKC with buffer D through a capillary with an inner coating of PEI. In the measurements, the reservoir that contained the oocyte **46'** was filled with buffer C, while the capillary **22**, the second reservoir **134** that served as the inlet for electrophoresis, and the outlet reservoir **34** contained buffer D.

Results of Measurements in Figs. 5 and 6A, 6B

Fig. 5 is an electrophoretogram of a sample of cytoplasm from an oocyte **46'** that was exposed to 10 μ M PMA for three minutes. Two fluorescent peaks, peak **192** corresponding to FI-sPKC and peak **194** corresponding to P-FI-sPKC, occur. This demonstrates that an intermediate degree of reporter substrate phosphorylation can be measured from a stimulated oocyte.

Fig. 6A is a graph of phosphorylation as a function of time showing the time course of intracellular phosphorylation of reporter substrate by PKC in response to extracellular treatment of *Xenopus laevis* oocytes with a potent pharmacologic specific activator of PKC. Fig. 6A shows the quantitative results of a number of measurements conducted to determine the time course of intracellular reporter substrate phosphorylation in oocytes **46'** in response to activation by exposure to 10 μ M PMA.

Fig. 6B is a graph of phosphorylation as a function of time showing the time course of intracellular phosphorylation of reporter substrate by PKC in response to extracellular treatment of *Xenopus laevis* oocytes with a physiological activator of PKC. Fig. 6B shows the quantitative results of a number of measurements conducted to determine the time course of

intracellular reporter substrate phosphorylation in oocytes 46' in response to activation by exposure to 10 μ M LPA. From the intermediate time points, FI-sPKC was phosphorylated within the oocytes 46' on average at a rate of 25 ± 15 nM/s as shown at a point 196 in Fig. 6A and 43 ± 15 nM/s as shown at a point 198 in Fig. 6B after exposure to PMA and LPA, respectively. At these rates, only $\sim 0.1\%$ of the peptide would have been phosphorylated in the 350 msec interval between the initiation of cytoplasmic sampling and electrophoresis.

Technical Aspects of the Measurement in Fig. 7

Fig. 7 is an electrophoretogram showing the separation of three different specific kinase reporter substrates from a sample of the contents of a *Xenopus laevis* oocyte that had been previously microinjected. In the measurement illustrated in Fig. 7, an individual oocyte 46' was microinjected with 50 nl of a mixture of reporter substrate peptides. The mixture was comprised of 20 μ M FI-sPKC, 7 μ M FI-sPKA, and 500 nM FI-scdc2K in buffer B. FI-sPKA, a specific reporter for protein kinase A (PKA) activity has the sequence FI-Lys-Arg-Arg-Glu-Ile-Leu-Ser-Arg-Arg-Pro-Ser-Tyr-Arg, and was derived from the CREB protein. FI-scdc2K, a specific reporter for cdc2 kinase (originally identified genetically as cell division cycle mutant 2) has the sequence FI-Gly-Gly-Gly-Arg-Ser-Pro-Gly-Arg-Arg-Arg-Lys, and comprises a consensus phosphorylation site derived from several proteins. The underlined serine residues are the sites of phosphorylation. The peptides were synthesized and labeled with fluorescein as described for FI-sPKC, except that FI-scdc2K was labeled with the mixed 5- and 6-isomers of carboxyfluorescein succinimidyl ester (100-200 mg/ml, Molecular Probes, Eugene, OR); thus, FI-scdc2k consisted of two isomeric forms. Each oocyte 46' was incubated in buffer C for 30 minutes to allow diffusion of the reporter substrates. The estimated final intracellular concentrations of reporter substrates were ~ 1 μ M FI-sPKC, ~ 330 nM FI-sPKA, and ~ 10 nM FI-scdc2K. Separation was achieved with PERT CE sampling followed by MEKC with buffer D through a capillary with an inner coating of PEI. In the measurement,

the reservoir that contained the oocyte **46'** was filled with buffer C, while the capillary **22**, the second reservoir **134**, and the outlet reservoir **34** contained buffer D.

Results of the Measurement in Fig. 7

5 Fig. 7 shows the separation of three different specific kinase reporter substrates from a sample of the contents of a *Xenopus laevis* oocyte. The peaks were identified by their migration times as observed when injected into oocytes **46'** singly (not shown). The first doublet, peaks **162** and **164**, corresponds to the two isomeric forms of nonphosphorylated FI-scdc2K. The
10 second doublet, peaks **166** and **168**, corresponds to the two isomeric forms of phosphorylated FI-scdc2K. Peak **170** is nonphosphorylated FI-sPKC and peak **172** is nonphosphorylated FI-sPKA. This measurement demonstrates that the net activities of multiple kinases can be measured from a single oocyte **46'**. Further, an exquisite degree of separation is achieved by MEKC
15 in this case as evidenced by the observable separation of isomeric forms of FI-scdc2K and P-FI-scdc2K. Finally, this measurement indicates that a substantial degree of cdc2K activity is present in an unstimulated oocyte **46'**, whereas PKC and PKA remain relatively inactive.

Many alterations and modifications may be made by those having
20 ordinary skill in the art without departing from the spirit and scope of the invention. Therefore, it must be understood that the illustrated embodiments have been set forth only for the purposes of example and that these should not be taken as limiting the invention as defined by the following claims.

The words used in this specification to describe the invention and its
25 various embodiments are to be understood not only in the sense of their commonly defined meanings, but to include by special definition in this specification structure, material or acts beyond the scope of the commonly defined meanings. Thus if an element can be understood in the context of this specification as including more than one meaning, then its use in a claim
30 must be understood as being generic to all possible meanings supported by the specification and by the word itself.

The definitions of the words or elements of the following claims are, therefore, defined in this specification to include not only the combination of elements which are literally set forth, but all equivalent structure, material or acts for performing substantially the same function in substantially the same way to obtain substantially the same result. In this sense it is therefore contemplated that an equivalent substitution of two or more elements may be made for any one of the elements in the claims below or that a single element may be substituted for two or more elements in a claim.

Insubstantial changes from the claimed subject matter as viewed by a person with ordinary skill in the art, now known or later devised, are expressly contemplated as being equivalently within the scope of the claims. Therefore, obvious substitutions now or later known to one with ordinary skill in the art are defined to be within the scope of the defined elements.

The claims are thus to be understood to include what is specifically illustrated and described above, what is conceptually equivalent, what can be obviously substituted and also what essentially incorporates the essential idea of the invention.

We claim:

1. A method for measuring the activity of intracellular chemical reactions in a cell comprising:
providing substrate molecules containing a label, the labeled
5 substrate molecules specific to a chemical reaction whose activity is to be measured;
disposing said substrate molecules within a single cell;
allowing said substrate molecules within the single cell to take part in the chemical reaction to produce altered substrate
10 molecules;
liberating said substrate molecules and said altered substrate molecules from the single cell;
detecting the label to identify the substrate molecules and the altered substrate molecules from the single cell; and
15 determining activity of said chemical reaction from a comparison of detected altered substrate molecules with detected substrate molecules.
2. The method of claim 1 further comprising quantifying the amounts of detected altered substrate molecules and detected substrate
20 molecules.
3. The method of claim 1 wherein said intracellular chemical reaction in said cell comprises enzyme catalysis.
4. The method of claim 1 wherein said altered substrate molecules exhibit a change in electrophoretic mobility as compared with the
25 substrate molecules.

5. The method of claim 4, wherein detecting said substrate molecules and said altered substrate molecules comprises electrophoresis.

6. The method of claim 2 wherein quantifying the amounts of detected altered substrate molecules and detected substrate molecules
5 comprises detection of the label by fluorescence following separation by electrophoresis.

7. The method of claim 1 wherein disposing said substrate molecules within a single cell comprises using a naturally occurring substrate molecule within said cell, inducing said substrate molecule to be produced within
10 said cell, or introducing said substrate molecule into said cell from outside said cell.

8. The method of claim 7 wherein introducing said substrate molecules into said cell from outside said cell comprises microinjecting, electroporating, optoporating, vesicle fusing, pinocytic loading, or associating
15 said substrate molecules with membrane permeant peptides.

9. The method of claim 1 further comprising stimulating said cell while said substrate molecules are intracellularly present prior to liberating said substrate molecules and said altered substrate molecules from the single cell.

20 10. The method of claim 9 further comprising comparing activity of said chemical reaction with a similar activity determined from a single cell that has not been stimulated.

11. The method of claim 1, wherein liberating said substrate molecules and said altered substrate molecules from the single cell comprises chemical disruption of said single cell, mechanical disruption of said single cell, or separation of said substrate and altered substrate molecules by
5 electrophoresis, or by a combination thereof.

12. The method of claim 1, wherein the label is selected from a group consisting of fluorescent labels, isotopes, labels exhibiting optical absorption, and electron spin resonance labels.

13. The method of claim 1 wherein the substrate molecules are
10 polymers.

14. The method of claim 13 wherein the polymers are selected from a group consisting of peptides, polysaccharide, and nucleic acids.

15. The method of claim 14 wherein said polymers are modified with a fluorescent label.

16. The method of claim 14 wherein said peptides are
15 substrates for a kinase that alters said modified peptides by the addition of a phosphate moiety to a particular amino acid within each peptide.

17. The method of claim 16, wherein said peptide has been modified by covalent addition of a fluorescent group.

18. The method of claim 1, said substrate molecules comprise
20 carbohydrates, phospholipids, entire proteins, or organic compounds not ordinarily found within the cell.

19. The method of claim 1 wherein detecting the label comprises performing voltammetry or mass spectrometry.

20. The method of claim 1 further comprising simultaneously performing each of said steps with a plurality of different substrate molecules,
5 each reporting on a specific chemical reaction within a single cell.

21. A method for measuring activity of a chemical reaction in a minute volume of tens of pl or less comprising:
providing substrate molecules containing a label;
disposing said substrate molecule into said minute volume where
10 said chemical reactions occurs producing altered substrate molecules within said minute volume;
terminating said chemical reactions;
detecting the label to identify the substrate molecules and the altered substrate molecules to determine activity of the
15 chemical reaction.

22. The method of claim 21 further comprising quantifying changes in the amounts of the substrate molecules and the altered substrate molecules.

23. An apparatus for measuring an activity of chemical reactions
20 of intracellular molecules comprising:
means for disposing labeled substrate molecules into said cell to form labeled altered substrate molecules therein; and
means for detecting said substrate molecules and said altered substrate molecules from a single cell before any
25 substantial alteration of said substrate molecules and said altered substrate molecules has occurred.

24. The apparatus of claim 23 further comprising means for quantifying changes in the amounts of said substrate molecules and said altered substrate molecules.

25. The apparatus of claim 23 wherein said means for detecting
5 detects enzyme catalysis.

26. The apparatus of claim 23 wherein said means for detecting detects a change in electrophoretic mobility of said substrate molecules versus said altered substrate molecules.

27. The apparatus of claim 23 wherein said means for detecting
10 comprises capillary electrophoresis.

28. The apparatus of claim 24 wherein said means for quantifying changes in the amounts of said substrate molecules and said altered substrate molecules comprises means for quantifying fluorescence of said substrate molecules and said altered substrate molecules following separation
15 of said substrate molecules and said altered substrate molecules by capillary electrophoresis.

29. The apparatus of claim 23 wherein said means for disposing comprises means for inducing said substrate molecules to be produced within said cell, or means for introducing said substrate molecule into said cell from
20 outside said cell.

30. The apparatus of claim 29 wherein said means for introducing said substrate molecules into said cell from outside said cell comprises means for microinjecting, means for electroporating, means for optoporing, means for vesicle fusing, means for pinocytic loading, or means
5 for associating said substrate molecules with membrane permeant peptides.

31. The apparatus of claim 23 wherein said means for disposing said substrate molecules comprises means for providing said substrate molecules from naturally occurring compounds or synthetically derived compounds.

10 32. The apparatus of claim 23 further comprising means for stimulating said cell while said substrate molecules are present intracellularly prior to detecting said substrate molecules and said altered substrate molecules.

33. The apparatus of claim 32 wherein said means for detecting comprises means for obtaining the contents of said cell, and means for
15 separating part or all of said contents by capillary electrophoresis

34. The apparatus of claim 33, wherein said means for obtaining the contents comprises chemical means for disruption, physical means for disruption, separation of molecules by electrophoresis, or a combination thereof.

35. The apparatus of claim 23 wherein said means for disposing
20 provides a substrate molecules that are specific for a particular intracellular chemical reaction.

36. The apparatus of claim 35 wherein said means for disposing provides substrate molecules which are fluorescent.

37. The apparatus of claim 36 wherein said means for disposing provides substrate molecules which are modified peptides.

38. The apparatus of claim 37, said modified peptides are substrates for a kinase that alters said modified peptides by the addition of a
5 phosphate moiety to a particular amino acid within each said peptide.

39. The apparatus of claim 37 wherein said peptides have been modified by covalent addition of a fluorescent group to allow detection by fluorescence.

40. The apparatus of claim 23 wherein said means for disposing
10 provides substrate molecules comprising nucleic acids, carbohydrates, phospholipids, entire proteins, or compounds not ordinarily found within cells.

41. The apparatus of claim 23 wherein said means for detecting comprises means for performing voltammetry or means for mass spectrometry on said substrate molecules and said altered substrate molecules.

15 42. The apparatus of claim 23 further comprising means for simultaneously disposing and detecting a plurality of different substrate molecules, each different substrate molecule reporting on a specific chemical reaction.

43. An apparatus for measuring an activity of chemical reactions
20 of molecules in a minute volume of the order of tens of pl or less comprising:
means for disposing substrate molecules having a label into said minute volume for chemical reactions to occur producing altered substrate molecules; and
means for detecting the label to identify the substrate molecules

and the altered substrate molecules to determine activity of the chemical reaction.

44. The apparatus of claim 43 further comprising means for quantifying changes in the amounts of said substrate molecules and said altered
5 substrate molecules.

45. An apparatus for measuring an activity of intracellular chemical reactions of molecules in a cell in which labeled substrate molecules have been disposed to allow for an in vivo reaction wherein labeled altered substrate molecules are formed, comprising:

10 a detector of said labeled substrate molecules and said labeled altered substrate molecules; and
a cell lysing device communicating with said detector, which cell lysing device extracts said substrate and altered substrate molecules from said cell, and which cell lysing device
15 collects and transfers said substrate and altered substrate molecules into said detector before any substantial alteration occurs.

46. The apparatus of claim 45 further comprising a data processor coupled to said detector to quantify changes in the amounts of said
20 substrate molecules and said altered substrate molecules.


47. An apparatus for measuring an activity of intracellular chemical reactions of molecules in a cell in which labeled substrate molecules have been disposed to allow for an in vivo reaction wherein labeled altered substrate molecules are formed, comprising:
25 means for holding the cell;
an electrophoresis reservoir contiguous to but not in fluidic contact with said means for holding;


a sharpened electrophoresis capillary for puncturing the cell to
remove a cellular sample;

means for moving said electrophoresis capillary to puncture said
cell;

5 means for rapidly transitioning said capillary into contact with said
electrophoresis reservoir after removing said cellular
sample so that electrophoresis of said cellular sample
through said capillary will commence; and


10 a detector for detecting said labeled substrate molecules and said
labeled altered substrate molecules during or following
electrophoresis.






3/11

FIG. 3



4/11

FIG. 4



5/11

FIG. 5




FIG. 6B






FIG. 6A



7/11



8/11

FIG. 8A






FIG. 8B



9/11

FIG. 9



10/11

FIG. 10






FIG. 12



11/11

FIG. 11



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/20009

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : G01N 33/48 US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 436/63, 164, 172; 422/82.05, 82.08; 435/4.29, 30, 288.7; 204/450, 451, 452, 453, 400, 403, 600, 601, 603, 604, 194 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WEST/USPAT, JPO, EPO, DERWENT; STN/CA, BIOSIS, MEDLINE search terms: enzyme, kinase, activity, single cell, capillary electrophoresis, fluorescence		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	LUZZI et al. Localized Sampling of Cytoplasm from Xenopus Oocytes for Capillary Electrophoresis. Analytical Chemistry. 1997, Vol. 69, pages 4761-4767, especially pages 4762 and 4766.	1-7, 11-15, 18, 21-29, 31, 35-36, 40, 43-47 ----- 8-10, 16-17, 19-20, 30, 32-34, 37-39, 41-42
Y	GIULIANO et al. Fluorescent Protein Biosensors: Measurement of Molecular Dynamics in Living Cells. Annu. Rev. Biophys. Biomol. Struct. 1995, Vol. 24, pages 405-434, especially pages 412 and 427.	8-10, 20, 30, 32-34, 42
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "N" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family	
Date of the actual completion of the international search 27 SEPTEMBER 2000		Date of mailing of the international search report 18 OCT 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer MAUREEN WALLENHORST Telephone No. (703) 308-0661 DEBORAH THOMAS PARALEGAL SPECIALIST

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/20009

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	HIGASHI et al. Imaging of cAMP-Dependent Protein Kinase Activity in Living Neural Cells Using a Novel Fluorescent Substrate. FEBS Letters. 1997, Vol. 414, pages 55-60, especially page 55.	16-17, 37-39
Y	LILLARD et al. Capillary Electrophoresis for the Analysis of Single Cells: Laser-Induced Fluorescence Detection. Handbook of Capillary Electrophoresis. 1997, J.P. Landers, Boca Raton, FL, CRC Press, pages 523-544, especially page 524.	19, 41
A	ZHANG et al. Detection and Differentiation of Pectic Enzyme Activity in vitro and in vivo by Capillary Electrophoresis of Products from Fluorescent-Labeled Substrate. Electrophoresis. 1996, Vol. 17, pages 372-378.	1-47
A	WU et al. Analysis of Src Kinase and Protein Kinase C Activity by Capillary Electrophoresis and Laser-Induced Fluorescence. Analytical Biochemistry. 1999, Vol. 269, pages 423-425.	1-47
A	SNOW et al. Determination of B1,4-Galactosyltransferase Enzymatic Activity by Capillary Electrophoresis and Laser-Induced Fluorescence Detection. Analytical Biochemistry. 1999, Vol. 271, pages 36-42.	1-47

INTERNATIONAL SEARCH REPORT

I. International application No.
PCT/US00/20009

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

436/63, 164, 172; 422/82.05, 82.08; 435/4.29, 288.7; 204/451, 452, 453, 403, 601, 603, 604

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.